


AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

Claim 1 (Previously Presented): A compound represented by the following formula (I) or a physiologically acceptable salt thereof, or a hydrate thereof:



wherein, R^1 and R^2 each independently represent hydrogen atom, a halogen atom, hydroxyl group, a group of OZ_{1-6} (the group of OZ_{1-6} represents an alkyl group having 1-6 carbon atoms or a fluoroalkyl group having 1-6 carbon atoms, which bonds via the oxygen atom), a group of $S(O)_nZ_{1-4}$ (Z_{1-4} represents an alkyl group having 1-4 carbon atoms or a fluoroalkyl group having 1-4 carbon atoms or an alkylene group derived therefrom), a group of $N(R^{12})(R^{13})$ (R^{12} and R^{13} each independently represent hydrogen atom, an alkyl group having 1-4 carbon atoms or a fluoroalkyl group having 1-4 carbon atoms), a group of Z_{1-8} which may be substituted (Z_{1-8} represents an alkyl group having 1-8 carbon atoms or a fluoroalkyl group having 1-8 carbon atoms), a 5- to 7-membered cyclic alkyl group, an aryl group, a heteroaryl group, or a 4- to 7-membered saturated or

partially saturated heterocyclic group (the cyclic alkyl group, aryl group, heteroaryl group and heterocyclic group may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a group of OZ_{1-4} , a group of $S(O)_nZ_{1-4}$, a group of $N(R^{12})(R^{13})$, a group of Z_{1-4} , carboxyl group, a group of CO_2Z_{1-4} , group of $CONH_2$, a group of $CONH(Z_{1-4})$ and a group of $CON(Z_{1-4})(Z_{1-4})$);

W^1 represents a group selected from the group consisting of $-CH=CH-$, $-N(R^{12})CO-$, $-CON(R^{12})-$, $-CH_2O-$ and $-CH_2CH_2-$ (each of the aforementioned groups binds to the thiazole ring at the left end);

R^3 represents hydrogen atom, a halogen atom, hydroxyl group or an amino group;

R^4 represents a group selected from the group consisting of hydrogen atom, a group of $-OZ_{0-4}R^5$ (Z_{0-4} represents an alkylene group having 1-4 carbon atoms, a fluorine-substituted alkylene group having 1-4 carbon atoms or a single bond, and R^5

represents a 5- to 7-membered cyclic alkyl group, an aryl group, a heteroaryl group or a 4- to 7-membered saturated or partially saturated heterocyclic group (the cyclic alkyl group, aryl group, heteroaryl group and heterocyclic group may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a group of OZ_{1-4} , a group of $S(O)_nZ_{1-4}$, a group of $N(R^{12})(R^{13})$, a group of Z_{1-4} , carboxyl group, a group of CO_2Z_{1-4} , group of $CONH_2$, a group of $CONH(Z_{1-4})$ and a group of $CON(Z_{1-4})(Z_{1-4})$), a group of $-S(O)_nZ_{0-4}R^5$, a group of $-N(R^6)(R^7)$ (R^6 and R^7 each independently represent hydrogen atom or Z_{1-4} , or they may bind to each other to form a saturated or unsaturated 5- to 7-membered ring (the ring may contain one or two hetero atoms as ring constituting atoms), and R^6 and R^7 may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a

group of $\text{OCON}(\text{R}^{12})(\text{R}^{13})$, a group of $\text{CON}(\text{R}^{12})(\text{R}^{13})$, a group of $\text{N}(\text{R}^{12})\text{CON}(\text{R}^{12})(\text{R}^{13})$, a group of Z_{1-4} , a group of OZ_{1-4} , a group $\text{S}(\text{O})_n\text{Z}_{1-4}$, group of CH_2OH , a group of $(\text{CH}_2)_m\text{N}(\text{R}^{12})(\text{R}^{13})$, carboxyl group, cyano group, a group of $\text{CO-Z}_{1-4}(\text{R}^{10})\text{-N}(\text{R}^{12})(\text{R}^{13})$ (R^{10} is a substituent corresponding to a side chain on an amino acid carbon or a group of $\text{-Z}_{1-4}\text{-R}^{11}$ (R^{11} represents a substituent which forms a quaternary salt) and a group of $\text{CO-Z}_{1-4}\text{-N}(\text{R}^{12})(\text{R}^{13})$ $\left. \begin{array}{c} | \\ (\text{CH}_2)_q \end{array} \right\}$, a 5- or 6-membered aryl group which may be substituted and a 5- or 6-membered unsaturated heterocyclic group which may be substituted;


W^2 represents a single bond or $\text{-C}(\text{R}^8)=\text{C}(\text{R}^9)\text{-}$ (R^8 and R^9 each independently represent hydrogen atom, a halogen atom, a lower alkyl group, an alkoxy group, cyano group, carboxyl group, hydroxymethyl group, cyanomethyl group, vinyl group or a group of $\text{N}(\text{R}^{12})(\text{R}^{13})$), Q represents an acidic group, and W^2 and Q may bind together to form vinylidenethiazolidinedione in *E*- or *Z*-configuration or an equivalent heterocyclic ring; m and n each independently represent an integer of 0 to 2, and q represents an integer of 0 to 3.

Claim 2 (Previously Presented): A medicament composition for eliminating resistance of a microorganism with acquired drug resistance, which comprises a compound represented by formula (I) according to claim 1 or a physiologically acceptable salt thereof as an active ingredient.

Claim 3 (Previously Presented): A medicament composition for enhancing effect of an antimicrobial agent, which comprises a compound represented by formula (I) according to claim 1 or a physiologically acceptable salt thereof as an active ingredient.

Claim 4 (Currently Amended): A pharmaceutical composition for preventive and/or therapeutic treatment of a microbial infection by microorganisms selected from *Pseudomonas aeruginosa* and bacteria having a genetically homologous drug efflux pump[,] which comprises a compound represented by formula (I) according to claim 1 or a physiologically acceptable salt thereof together with an antimicrobial agent.

Claim 5 (Previously Presented): A compound represented by the following formula (I) or a physiologically acceptable salt thereof, or hydrate thereof



wherein, R^1 and R^2 each independently represent hydrogen atom, a halogen atom, hydroxyl group, a group of OZ_{1-6} (the group of OZ_{1-6} represents an alkyl group having 1-6 carbon atoms or a fluoroalkyl group having 1-6 carbon atoms, which bonds via the oxygen atom), a group of $S(O)_nZ_{1-4}$ (Z_{1-4} represents an alkyl group having 1-4 carbon atoms or a fluoroalkyl group having 1-4 carbon atoms or an alkylene group derived therefrom), a group of $N(R^{12})(R^{13})$ (R^{12} and R^{13} each independently represent hydrogen atom, an alkyl group having 1-4 carbon atoms or a fluoroalkyl group having 1-4 carbon atoms), a group of Z_{1-8} which may be substituted (Z_{1-8} represents an alkyl group having 1-8 carbon atoms or a fluoroalkyl group having 1-8 carbon atoms), a 5- to 7-membered cyclic alkyl group, an aryl group, a heteroaryl group, or a 4- to 7-membered saturated or partially saturated heterocyclic group (the cyclic alkyl group, aryl group, heteroaryl group and heterocyclic group may have one to three substituents selected from the group

consisting of a halogen atom, hydroxyl group, a group of OZ_{1-4} , a group of $S(O)_nZ_{1-4}$, a group of $N(R^{12})(R^{13})$, a group of Z_{1-4} , carboxyl group, a group of CO_2Z_{1-4} , group of $CONH_2$, a group of $CONH(Z_{1-4})$ and a group of $CON(Z_{1-4})(Z_{1-4})$;

W^1 represents a group selected from the group consisting of $-CH=CH-$, $-N(R^{12})CO-$, $-CON(R^{12})-$, $-CH_2O-$ and $-CH_2CH_2-$ (each of the aforementioned groups binds to the thiazole ring at the left end);

R^3 represents hydrogen atom, a halogen atom, hydroxyl group or an amino group;

R^4 represents a group selected from the group consisting of hydrogen atom, a group of $-OZ_{0-4}R^5$ (Z_{0-4} represents an alkylene group having 1-4 carbon atoms, a fluorine-substituted alkylene group having 1-4 carbon atoms or a single bond, and R^5 represents a 5- to 7-membered cyclic alkyl group, an aryl group, a heteroaryl group or a 4- to 7-membered saturated or partially saturated heterocyclic group (the cyclic alkyl group, aryl group, heteroaryl group and heterocyclic group may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a group of OZ_{1-4} , a group of $S(O)_nZ_{1-4}$, a group of $N(R^{12})(R^{13})$, a group of Z_{1-4} , carboxyl group, a group of CO_2Z_{1-4} , group of $CONH_2$, a group of $CONH(Z_{1-4})$ and a group of $CON(Z_{1-4})(Z_{1-4})$), a group of $-S(O)_nZ_{0-4}R^5$, a group of $-N(R^6)(R^7)$ (R^6 and R^7 each independently represent hydrogen atom or Z_{1-4} , or they may bind to each other to form a saturated or unsaturated 5- to 7-membered ring (the ring may contain one or two hetero atoms as ring constituting atoms), and R^6 and R^7 may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a group of $OCON(R^{12})(R^{13})$, a group of $CON(R^{12})(R^{13})$, a group of $N(R^{12})CON(R^{12})(R^{13})$, a group of Z_{1-4} , a group of OZ_{1-4} , a group $S(O)_nZ_{1-4}$, group of CH_2OH , a group of

$(CH_2)_m N(R^{12})(R^{13})$, carboxyl group, cyano group, a group of $CO-Z_{1-4}(R^{10})-N(R^{12})(R^{13})$
 $(R^{10}$ is a substituent corresponding to a side chain on an amino acid carbon or a group
of $-Z_{1-4}-R^{11}$ (R^{11} represents a substituent which forms a quaternary salt) and a group of
 $CO-Z_{1-4}-N(R^{12})(R^{13})$
 $(CH_2)_q$ }, a 5- or 6-membered aryl group which may be substituted and a 5-
or 6-membered unsaturated heterocyclic group which may be substituted;
 W^2 represents a single bond or $-C(R^8)=C(R^9)-$ (R^8 and R^9 each independently represent
hydrogen atom, a halogen atom, a lower alkyl group, an alkoxy group, cyano group,
carboxyl group, hydroxymethyl group, cyanomethyl group, vinyl group or a group of
 $N(R^{12})(R^{13})$), Q represents an acidic group, and W^2 and Q may bind together to form
vinylidenethiazolidinedione in *E*- or *Z*-configuration or an equivalent heterocyclic ring; m
and n each independently represent an integer of 0 to 2, and q represents an integer of
0 to 3; R^{14} represents hydrogen atom, Z_{1-4} , $Z_{1-4}R^5$ or
 $Z_{1-4}OR^5$; and X represents C-H and Y represents C-H or nitrogen atom.

Claim 6 (Currently Amended): A medicament composition for ~~preventive and/or~~
therapeutic treatment of ~~a microbial infection~~ by microorganisms selected from
Pseudomonas aeruginosa and bacteria having a genetically homologous drug efflux
pump which comprises a compound represented by the formula (I) according to claim 1
or a physiologically acceptable salt thereof as an active ingredient.

Claim 7 (Currently Amended): A method for therapeutic treatment of ~~a microbial~~
infection by microorganisms selected from *Pseudomonas aeruginosa* and bacteria
having a genetically homologous drug efflux pump comprising administering to a
mammal in need thereof a therapeutically effective amount of the composition

according to claim 6.

Claim 8 (Previously Presented): The method according to claim 7, further comprising administering at least one antimicrobial agent.

Claim 9 (Previously Presented) The method according to claim 8, wherein the at least one antimicrobial agent is simultaneously administered with the composition.

Claim 10 (Previously Presented): The method according to claim 8, wherein the at least one antimicrobial agent is separately administered from the composition.

Claim 11 (Previously Presented): The method according to claim 8, wherein the at least one antimicrobial agent is successively administered with the composition.

Claim 12 (Previously Presented): The method according to claim 7 wherein the mammal is a human.

Claim 13 (Previously Presented): A method for preventive treatment of a microbial infection comprising administering to a mammal a preventively effective amount of the composition according to claim 6.

Claim 14 (Previously Presented): The method according to claim 13, further comprising administering at least one antimicrobial agent.

Claim 15 (Previously Presented): The method according to claim 14, wherein the at least one antimicrobial agent is simultaneously administered with the composition.

Claim 16 (Previously Presented): The method according to claim 14, wherein the at least one antimicrobial agent is separately administered from the composition.

Claim 17 (Previously Presented): The method according to claim 14, wherein the at least one antimicrobial agent is successively administered with the composition.

Claim 18 (Previously Presented): The method according to claim 13 wherein the mammal is a human.

Claim 19 (Canceled)

Claim 20 (Currently Amended): A medicament composition for ~~preventive and/or~~ therapeutic treatment of ~~a microbial-infection~~ by microorganisms selected from Pseudomonas aeruginosa and bacteria having a genetically homologous drug efflux pump which comprises a compound represented by the formula (I) according to claim 5 or a physiologically acceptable salt thereof as an active ingredient.


Claim 21 (Canceled)

Claim 22 (Currently Amended): A method for therapeutic treatment of a ~~microbial-infection~~ by microorganisms selected from Pseudomonas aeruginosa and bacteria having a genetically homologous drug efflux pump comprising administering to a mammal in need thereof a therapeutically effective amount of the composition according to claim 20.

Claim 23 (Canceled)

Claim 24 (Previously Presented): The method according to claim 22, further comprising administering at least one antimicrobial agent.

Claim 25. (Currently Amended): A method for therapeutic treatment of a ~~microbial-infection~~ by microorganisms selected from Pseudomonas aeruginosa and bacteria having a genetically homologous drug efflux pump comprising administering to a mammal in need thereof a therapeutically effective amount of a composition comprising a compound represented by formula (I) or a physiologically acceptable salt thereof as an active ingredient and at least one antimicrobial agent



wherein, R^1 and R^2 each independently represent hydrogen atom, a halogen atom, hydroxyl group, a group of OZ_{1-6} (the group of OZ_{1-6} represents an alkyl group having 1-6 carbon atoms or a fluoroalkyl group having 1-6 carbon atoms, which bonds via the oxygen atom), a group of $S(O)_nZ_{1-4}$ (Z_{1-4} represents an alkyl group having 1-4 carbon atoms or a fluoroalkyl group having 1-4 carbon atoms or an alkylene group derived therefrom), a group of $N(R^{12})(R^{13})$ (R^{12} and R^{13} each independently represent hydrogen atom, an alkyl group having 1-4 carbon atoms or a fluoroalkyl group having 1-4 carbon atoms), a group of Z_{1-8} which may be substituted (Z_{1-8} represents an alkyl group having 1-8 carbon atoms or a fluoroalkyl group having 1-8 carbon atoms), a 5- to 7-membered cyclic alkyl group, an aryl group, a heteroaryl group, or a 4- to 7-membered saturated or partially saturated heterocyclic group (the cyclic alkyl group, aryl group, heteroaryl group and heterocyclic group may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a group of OZ_{1-4} , a group of $S(O)_nZ_{1-4}$, a group of $N(R^{12})(R^{13})$, a group of Z_{1-4} , carboxyl group, a group of CO_2Z_{1-4} , group of $CONH_2$, a group of $CONH(Z_{1-4})$ and a group of $CON(Z_{1-4})(Z_{1-4})$);

W^1 represents a group selected from the group consisting of $-CH=CH-$, $-N(R^{12})CO-$, $-CON(R^{12})-$, $-CH_2O-$ and $-CH_2CH_2-$ (each of the aforementioned groups binds to the thiazole ring at the left end);

R^3 represents hydrogen atom, a halogen atom, hydroxyl group or an amino group;


R^4 represents a group selected from the group consisting of hydrogen atom, a group of $-OZ_{0-4}R^5$ (Z_{0-4} represents an alkylene group having 1-4 carbon atoms, a fluorine-substituted alkylene group having 1-4 carbon atoms or a single bond, and R^5 represents a 5- to 7-membered cyclic alkyl group, an aryl group, a heteroaryl group or a 4- to 7-membered saturated or partially saturated heterocyclic group (the cyclic alkyl group, aryl group, heteroaryl group and heterocyclic group may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a group of OZ_{1-4} , a group of $S(O)_nZ_{1-4}$, a group of $N(R^{12})(R^{13})$, a group of Z_{1-4} , carboxyl group, a group of CO_2Z_{1-4} , group of $CONH_2$, a group of $CONH(Z_{1-4})$ and a group of $CON(Z_{1-4})(Z_{1-4})$), a group of $-S(O)_nZ_{0-4}R^5$, a group of $-N(R^6)(R^7)$ (R^6 and R^7 each independently represent hydrogen atom or Z_{1-4} , or they may bind to each other to form a saturated or unsaturated 5- to 7-membered ring (the ring may contain one or two hetero atoms as ring constituting atoms), and R^6 and R^7 may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a group of $OCON(R^{12})(R^{13})$, a group of $CON(R^{12})(R^{13})$, a group of $N(R^{12})CON(R^{12})(R^{13})$, a group of Z_{1-4} , a group of OZ_{1-4} , a group $S(O)_nZ_{1-4}$, group of CH_2OH , a group of $(CH_2)_mN(R^{12})(R^{13})$, carboxyl group, cyano group, a group of $CO-Z_{1-4}(R^{10})-N(R^{12})(R^{13})$ (R^{10} is a substituent corresponding to a side chain on an amino acid carbon or a group of $-Z_{1-4}-R^{11}$ (R^{11} represents a substituent which forms a quaternary salt) and a group of $CO-Z_{1-4}-N(R^{12})(R^{13})$), a 5- or 6-membered aryl group which may be substituted and a 5- or 6-membered unsaturated heterocyclic group which may be substituted;

W^2 represents a single bond or $-C(R^8)=C(R^9)-$ (R^8 and R^9 each independently represent

hydrogen atom, a halogen atom, a lower alkyl group, an alkoxy group, cyano group, carboxyl group, hydroxymethyl group, cyanomethyl group, vinyl group or a group of $N(R^{12})(R^{13})$, Q represents an acidic group, and W^2 and Q may bind together to form vinylidenethiazolidinedione in *E*- or *Z*-configuration or an equivalent heterocyclic ring; m and n each independently represent an integer of 0 to 2, and q represents an integer of 0 to 3; R^{14} represents hydrogen atom, an alkyl group having 1, 3 or 4 carbon atoms or a fluoroalkyl group having 1-4 carbon atoms, $Z_{1-4}R^5$ or $Z_{1-4}OR^5$; and X and Y each independently represent C-H or nitrogen atom.

Claim 26 (Currently Amended): A method for preventive treatment of a microbial infection by microorganisms selected from *Pseudomonas aeruginosa* and bacteria having a genetically homologous drug efflux pump comprising administering to a mammal a preventively effective amount of the composition according to claim 20.

Claim 27 (Currently Amended): A method for preventive treatment of a microbial infection by microorganisms selected from *Pseudomonas aeruginosa* and bacteria having a genetically homologous drug efflux pump comprising administering to a mammal a preventively effective amount of a composition comprising a compound represented by the formula (I) or a physiologically acceptable salt thereof as an active ingredient and at least one antimicrobial agent[.]



(I)

wherein, R^1 and R^2 each independently represent hydrogen atom, a halogen atom,

hydroxyl group, a group of OZ_{1-6} (the group of OZ_{1-6} represents an alkyl group having 1-6 carbon atoms or a fluoroalkyl group having 1-6 carbon atoms, which bonds via the oxygen atom), a group of $S(O)_nZ_{1-4}$ (Z_{1-4} represents an alkyl group having 1-4 carbon atoms or a fluoroalkyl group having 1-4 carbon atoms or an alkylene group derived therefrom), a group of $N(R^{12})(R^{13})$ (R^{12} and R^{13} each independently represent hydrogen atom, an alkyl group having 1-4 carbon atoms or a fluoroalkyl group having 1-4 carbon atoms), a group of Z_{1-8} which may be substituted (Z_{1-8} represents an alkyl group having 1-8 carbon atoms or a fluoroalkyl group having 1-8 carbon atoms), a 5- to 7-membered cyclic alkyl group, an aryl group, a heteroaryl group, or a 4- to 7-membered saturated or partially saturated heterocyclic group (the cyclic alkyl group, aryl group, heteroaryl group and heterocyclic group may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a group of OZ_{1-4} , a group of $S(O)_nZ_{1-4}$, a group of $N(R^{12})(R^{13})$, a group of Z_{1-4} , carboxyl group, a group of CO_2Z_{1-4} , group of $CONH_2$, a group of $CONH(Z_{1-4})$ and a group of $CON(Z_{1-4})(Z_{1-4})$);

W^1 represents a group selected from the group consisting of $-CH=CH-$, $-N(R^{12})CO-$, $-CON(R^{12})-$, $-CH_2O-$ and $-CH_2CH_2-$ (each of the aforementioned groups binds to the thiazole ring at the left end);

R^3 represents hydrogen atom, a halogen atom, hydroxyl group or an amino group;

R^4 represents a group selected from the group consisting of hydrogen atom, a group of $-OZ_{0-4}R^5$ (Z_{0-4} represents an alkylene group having 1-4 carbon atoms, a fluorine-substituted alkylene group having 1-4 carbon atoms or a single bond, and R^5 represents a 5- to 7-membered cyclic alkyl group, an aryl group, a heteroaryl group or a 4- to 7-membered saturated or partially saturated heterocyclic group (the cyclic alkyl

group, aryl group, heteroaryl group and heterocyclic group may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a group of OZ_{1-4} , a group of $\text{S(O)}_n\text{Z}_{1-4}$, a group of $\text{N(R}^{12})(\text{R}^{13})$, a group of Z_{1-4} , carboxyl group, a group of $\text{CO}_2\text{Z}_{1-4}$, group of CONH_2 , a group of $\text{CONH(Z}_{1-4})$ and a group of $\text{CON(Z}_{1-4})(\text{Z}_{1-4})$, a group of $-\text{S(O)}_n\text{Z}_{0-4}\text{R}^5$, a group of $-\text{N(R}^6)(\text{R}^7)$ (R^6 and R^7 each independently represent hydrogen atom or Z_{1-4} , or they may bind to each other to form a saturated or unsaturated 5- to 7-membered ring (the ring may contain one or two hetero atoms as ring constituting atoms), and R^6 and R^7 may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a group of $\text{OCON(R}^{12})(\text{R}^{13})$, a group of $\text{CON(R}^{12})(\text{R}^{13})$, a group of $\text{N(R}^{12})\text{CON(R}^{12})(\text{R}^{13})$, a group of Z_{1-4} , a group of OZ_{1-4} , a group $\text{S(O)}_n\text{Z}_{1-4}$, group of CH_2OH , a group of $(\text{CH}_2)_m\text{N(R}^{12})(\text{R}^{13})$, carboxyl group, cyano group, a group of $\text{CO-Z}_{1-4}(\text{R}^{10})-\text{N(R}^{12})(\text{R}^{13})$ (R^{10} is a substituent corresponding to a side chain on an amino acid carbon or a group of $-\text{Z}_{1-4}-\text{R}^{11}$ (R^{11} represents a substituent which forms a quaternary salt) and a group of $\text{CO-Z}_{1-4}-\text{N(R}^{12})(\text{R}^{13})$ $\left. \begin{array}{c} | \\ (\text{CH}_2)_q \end{array} \right\}$, a 5- or 6-membered aryl group which may be substituted and a 5- or 6-membered unsaturated heterocyclic group which may be substituted;

W^2 represents a single bond or $-\text{C(R}^8)=\text{C(R}^9)-$ (R^8 and R^9 each independently represent hydrogen atom, a halogen atom, a lower alkyl group, an alkoxy group, cyano group, carboxyl group, hydroxymethyl group, cyanomethyl group, vinyl group or a group of $\text{N(R}^{12})(\text{R}^{13})$), Q represents an acidic group, and W^2 and Q may bind together to form vinylidenethiazolidinedione in *E*- or *Z*-configuration or an equivalent heterocyclic ring; m and n each independently represent an integer of 0 to 2, and q represents an integer of 0

to 3; R¹⁴ represents hydrogen atom, an alkyl group having 1, 3 or 4 carbon atoms or a fluoroalkyl group having 1-4 carbon atoms, Z₁₋₄R⁵ or Z₁₋₄OR⁵; and X and Y each independently represent C-H or nitrogen atom.


Claim 28 (Currently Amended): A method for inhibiting drug resistance acquisition due to a drug efflux pump comprising administering to a mammal in need thereof an effective amount to inhibit drug resistance acquisition due to a drug efflux pump of the composition according to claim 6.

Claim 29 (Previously Presented): The method according to claim 28 wherein the mammal is a human.

Claim 30 (Currently Amended): A method for inhibiting drug resistance acquisition due to a drug efflux pump comprising administering to a mammal in need thereof an effective amount to inhibit drug resistance acquisition due to a drug efflux pump of the composition according to claim 20.

Claim 31 (Previously Presented): The method according to claim 30 wherein the mammal is a human.

Claim 32 (Currently Amended): A method for inhibiting drug resistance acquisition due to a drug efflux pump comprising administering to a mammal in need thereof an effective amount to inhibit drug resistance acquisition due to a drug efflux pump of a composition comprising a compound represented by formula (I) or a physiologically acceptable salt thereof as an active ingredient



wherein, R^1 and R^2 each independently represent hydrogen atom, a halogen atom, hydroxyl group, a group of OZ_{1-6} (the group of OZ_{1-6} represents an alkyl group having 1-6 carbon atoms or a fluoroalkyl group having 1-6 carbon atoms, which bonds via the oxygen atom), a group of $S(O)_nZ_{1-4}$ (Z_{1-4} represents an alkyl group having 1-4 carbon atoms or a fluoroalkyl group having 1-4 carbon atoms or an alkylene group derived therefrom), a group of $N(R^{12})(R^{13})$ (R^{12} and R^{13} each independently represent hydrogen atom, an alkyl group having 1-4 carbon atoms or a fluoroalkyl group having 1-4 carbon atoms), a group of Z_{1-8} which may be substituted (Z_{1-8} represents an alkyl group having 1-8 carbon atoms or a fluoroalkyl group having 1-8 carbon atoms), a 5- to 7-membered cyclic alkyl group, an aryl group, a heteroaryl group, or a 4- to 7-membered saturated or partially saturated heterocyclic group (the cyclic alkyl group, aryl group, heteroaryl group and heterocyclic group may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a group of OZ_{1-4} , a group of $S(O)_nZ_{1-4}$, a group of $N(R^{12})(R^{13})$, a group of Z_{1-4} , carboxyl group, a group of CO_2Z_{1-4} , group of $CONH_2$, a group of $CONH(Z_{1-4})$ and a group of $CON(Z_{1-4})(Z_{1-4})$);

W^1 represents a group selected from the group consisting of $-CH=CH-$, $-N(R^{12})CO-$, $-CON(R^{12})-$, $-CH_2O-$ and $-CH_2CH_2-$ (each of the aforementioned groups binds to the thiazole ring at the left end);

R^3 represents hydrogen atom, a halogen atom, hydroxyl group or an amino group;

R^4 represents a group selected from the group consisting of hydrogen atom, a group of $-OZ_{0-4}R^5$ (Z_{0-4} represents an alkylene group having 1-4 carbon atoms, a fluorine-substituted alkylene group having 1-4 carbon atoms or a single bond, and R^5 represents a 5- to 7-membered cyclic alkyl group, an aryl group, a heteroaryl group or a 4- to 7-membered saturated or partially saturated heterocyclic group (the cyclic alkyl group, aryl group, heteroaryl group and heterocyclic group may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a group of OZ_{1-4} , a group of $S(O)_nZ_{1-4}$, a group of $N(R^{12})(R^{13})$, a group of Z_{1-4} , carboxyl group, a group of CO_2Z_{1-4} , group of $CONH_2$, a group of $CONH(Z_{1-4})$ and a group of $CON(Z_{1-4})(Z_{1-4})$), a group of $-S(O)_nZ_{0-4}R^5$, a group of $-N(R^6)(R^7)$ (R^6 and R^7 each independently represent hydrogen atom or Z_{1-4} , or they may bind to each other to form a saturated or unsaturated 5- to 7-membered ring (the ring may contain one or two hetero atoms as ring constituting atoms), and R^6 and R^7 may have one to three substituents selected from the group consisting of a halogen atom, hydroxyl group, a group of $OCON(R^{12})(R^{13})$, a group of $CON(R^{12})(R^{13})$, a group of $N(R^{12})CON(R^{12})(R^{13})$, a group of Z_{1-4} , a group of OZ_{1-4} , a group $S(O)_nZ_{1-4}$, group of CH_2OH , a group of $(CH_2)_mN(R^{12})(R^{13})$, carboxyl group, cyano group, a group of $CO-Z_{1-4}(R^{10})-N(R^{12})(R^{13})$ (R^{10} is a substituent corresponding to a side chain on an amino acid carbon or a group of $-Z_{1-4}-R^{11}$ (R^{11} represents a substituent which forms a quaternary salt) and a group of $CO-Z_{1-4}-N(R^{12})(R^{13})$ $(CH_2)_q$), a 5- or 6-membered aryl group which may be substituted and a 5- or 6-membered unsaturated heterocyclic group which may be substituted;

W^2 represents a single bond or $-C(R^8)=C(R^9)-$ (R^8 and R^9 each independently

represent hydrogen atom, a halogen atom, a lower alkyl group, an alkoxy group, cyano group, carboxyl group, hydroxymethyl group, cyanomethyl group, vinyl group or a group of $N(R^{12})(R^{13})$, Q represents an acidic group, and W^2 and Q may bind together to form vinylidenethiazolidinedione in *E*- or *Z*-configuration or an equivalent heterocyclic ring; m and n each independently represent an integer of 0 to 2, and q represents an integer of 0 to 3; R^{14} represents hydrogen atom, an alkyl group having 1, 3 or 4 carbon atoms or a fluoroalkyl group having 1-4 carbon atoms, $Z_{1-4}R^5$ or $Z_{1-4}OR^5$; and X and Y each independently represent C-H or nitrogen atom.

33 (Previously Presented): The method according to claim 32 wherein the mammal is a human.